

Ascension Borgess Orthopedics & Sports Medicine

Postop Rehabilitation

Arthroscopic Superior Capsular Reconstruction (SCR)

Phase I (0-6 weeks)

Range of Motion: Limit FF 90°, ER 45°, Extension 20°, Abduction 45°, ABER 45°

Sling: Off for home exercises and hygiene.

Exercises: Elbow/wrist ROM. Pendulums, saws, table-slides 4x/d x 10min.

Restrictions: No lifting heavier than a coffee cup.

Phase II (6-12 weeks)

Range of Motion: Begin active/active assisted ROM. Advance to FF 140°, Abduction 135°, ABER 90°, ABIR 45°.

Sling: None

Exercises: Light deltoid/rotator cuff isometrics. Scapular stabilization. Light rotator cuff strengthening, biceps, triceps.

Restrictions: No aggressive stretching. No extended arm lifting.

Phase III (3-5 months)

Range of Motion: Gradual return to full AROM.

Sling: None

Exercises: Increase ER and latissimus eccentrics, glenohumeral stabilization. Begin rotator cuff endurance. Cycling/running as tolerated.

Phase IV (5-8 months)

Range of Motion: Full and pain-free

Sling: None

Exercises: Scapular stabilization, eccentric strengthening, and perturbation. Plyometric and throwing/racquet program. Continue endurance. Return to full activity as tolerated.

www.christopheruggenmd.com

2490 S. 11th St, Kalamazoo, MI 49009 • 269-343-1535